



Abstract

Aglycosylated IgG (including IgG1, IgG2 and IgG4 subclasses) were synthesized using PUREfrex. The PURE (Protein synthesis Using Recombinant Elements) system is a reconstituted cell-free protein synthesis system, which consists of only purified factors necessary for transcription, translation and energy regeneration. Recently, we developed an updated PURE system with higher productivity, which was launched as "PUREfrex 2.0" in 2015. We reported that PUREfrex 2.0 could be used for production of fragment antibodies such as Fab and scFv by last year. Here, we report the further application using PUREfrex 2.0 for production of IgG. We chose Trastuzumab as a model protein and optimized the composition of PUREfrex 2.0 reaction mixture and the reaction conditions as below; 1) adding disulfide bond isomerase (DsbC); 2) choosing GSH as an effective reductant; 3) adjusting GSH/GSSG ratio; 4) adding molecular chaperone (DnaK mix); 5) long-time incubation for 28h. We also suggested that synthesis temperature and template DNA ratio (light chain/heavy chain) should be optimized for individual IgGs for the best yield. At the best mode of synthesis, the productivity of Trastuzumab reached to 124 µg/mL. Moreover, the 68 µg of purified Trastuzumab was obtained from 1 mL of the reaction mixture after the purification by protein A resin and the following gel filtration. The purified Trastuzumab exhibited high binding affinity to recombinant HER2 protein (KD=4.24E-10 M) and internalized into HER2 expressing BT-474 cells. Furthermore, other IgGs including IgG1, IgG2 and IgG4 subclasses were also synthesized and confirmed their binding activity in ELISA. These results indicate that PUREfrex will be useful tool for high-throughput production/screening of functional antibodies (scFv, Fab, and IgG).


1. Optimization of the synthesis of Trastuzumab using PUREfrex®



2. Evaluation of the synthesized Trastuzumab



3. Synthesis of other IgGs using PUREfrex®



Method for the synthesis of IgG using PUREfrex

Reaction mix: **PUREfrex 2.0 Customized solution (without DTT and with 2 mM GSH)** + **DS supplement (3 mM GSSG, 5 µM (117 µg/mL) DsbC)** + **DnaK Mix (5 µM DnaK, 1 µM DnaJ, 1 µM GrpE)**

Incubation: **30 - 37°C for 28h**

Template DNA: **PCR product (final 10 nM mixed template DNA)**

Synthesis temperature and template DNA ratio (light chain/heavy chain) should be optimized for individual IgGs for the best yield.

*If you want to synthesize multiple IgGs at the same time (e.g. IgG screening), we recommend "30°C and LC:HC=1:1" as a first step.

Conclusion

- Aglycosylated IgG (including IgG1, IgG2 and IgG4 subclasses) was synthesized in the active form using PUREfrex 2.0.
- Disulfide bond isomerase (DsbC), redox state (reducing agent/oxidizing agent ratio) and long-time incubation were important to form a correct hetero tetramer of two heavy chains and two light chains.
- IgG yield, optimum synthesis temperature and optimum template DNA ratio were different between individual IgGs, which may reflect the difference of CDR sequence and structural stability.
- PUREfrex® will be "useful" and "powerful" tool for high-throughput production/screening of functional antibodies.

Acknowledgements

We thank Dr. Yosuke Uchinashi (Nihon Pall Ltd.) and Keita Iguchi (KANEKA Corporation) for technical support and experimental advice.